Clinical documentation after rapid response system calls	
IDENTIFYING AND DEFINITIONAL ATTRIBUTES	
SHORT NAME:	Clinical documentation after rapid response system calls
DESCRIPTION:	The proportion of rapid response system calls for which there is documentation in the clinical record of the details of the event
TYPE OF QUALITY MEASURE:	Process measure
RATIONALE:	Inadequate clinical documentation has been identified as an important contributing factor to adverse events in healthcare. Poor written and verbal communication between health professionals can result in discontinuity of care, delays in treatment, adverse events and increased morbidity and mortality. Poor communication also poses risks to patient safety when patients are transferred between clinical areas and during critical events such as rapid response system calls
DEFINITIONS:	Admitted patient: any patient for whom the hospital accepts responsibility for the provision of inpatient care and/or treatment. Admission follows a clinical decision based upon specified criteria that a patient requires same day or overnight care or treatment
	Evidence of clinical documentation: documentation in the healthcare record that summarises the details of the rapid response call and meets any requirements outlined in the facility's rapid response policy
	Rapid response system: system that provides emergency assistance to patients whose condition is deteriorating
	Rapid response system call: presence of either a rapid response system call record form in the patient's healthcare record or other relevant documentation
COLLECTION AND USAGE ATTRIBUTES	
POPULATION:	Admitted patients who receive a rapid response system call
COMPUTATION:	Percentage of rapid response system calls for which there is a documented summary of the details of the call in accordance with the requirements of rapid response policy Numerator × 100 Denominator
NUMERATOR:	Number of audited rapid response system calls for which there is a documented summary of the details of the call in accordance with the requirements of rapid response policy
DENOMINATOR:	Total number of audited rapid response system calls
COMMENTS	
COMMENTS:	A high rate of clinical documentation after rapid response calls is desirable
	Evidence of clinical documentation should be assessed in accordance with the agreed documentation process outlined in the facility's rapid response policy
	Collecting data for this quality measure will require information from the records of rapid response system calls and from the patient's healthcare record

Clinical documentation after rapid response system calls

REFERENCES

REFERENCE DOCUMENTS:

Australian Commission on Safety and Quality in Health Care. OSSIE Guide to Clinical Handover Improvement. Sydney. ACSQHC, 2009

Australian Institute of Health and Welfare Index. (Accessed 5 August 2011, at http://meteor.aihw.gov.au/content/index.phtml/itemId/327206.)

National Institute for Health and Clinical Excellence. Acutely ill patients in hospital: Recognition of and response to acute illness in adults in hospital. National Institute for Health and Clinical Excellence, 2007

Sebat F. Designing, implementing and enhancing a Rapid Response System. Mount Prospect: Society of Critical Care Medicine, 2009